

§ 640.22

(b) Plasmapheresis donors shall meet the criteria for suitability prescribed in § 640.63, excluding the phrase "other than malaria" in paragraph (c)(9). Informed consent shall be required as prescribed in § 640.61.

(c) Plateletpheresis donors shall meet criteria for suitability as described in a license application or a supplement to the product license, and must have the written approval of the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

[40 FR 4304, Jan. 29, 1975, as amended at 49 FR 23834, June 8, 1984; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994]

§ 640.22 Collection of source material.

(a) Whole blood used as the source of Platelets shall be collected as prescribed in § 640.4, except that paragraphs (d)(2) and (h) shall not apply.

(b) If plasmapheresis is used, the procedure for collection shall be prescribed in §§ 640.62, 640.64 (except paragraph (c)(3)), and 640.65.

(c) If plateletpheresis is used, the procedure for collection shall be as described in a license application or a supplement to a product license, and must have the written approval of the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

(d) The phlebotomy shall be performed by a single uninterrupted venipuncture with minimal damage to, and minimal manipulation of, the donor's tissue.

[40 FR 4304, Jan. 29, 1975, as amended at 45 FR 27927, Apr. 25, 1980; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994]

§ 640.23 Testing the blood.

(a) Blood from which plasma is separated for the preparation of Platelets shall be tested as prescribed in §§ 610.40 and 610.45 of this chapter and § 640.5 (a), (b), and (c).

(b) The tests shall be performed on a sample of blood collected at the time of collecting the source blood, and such sample container shall be labeled with

21 CFR Ch. I (4–1–98 Edition)

the donor's number before the container is filled.

[40 FR 4304, Jan. 29, 1975, as amended at 50 FR 4139, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988]

§ 640.24 Processing.

(a) Separation of plasma and platelets and resuspension of the platelets shall be in a closed system. Platelets shall not be pooled during processing.

(b) Immediately after collection, the whole blood or plasma shall be held in storage between 20 to 24 °C, unless it must be transported from the donor clinic to the processing laboratory. During such transport, all reasonable methods shall be used to maintain the temperature as close as possible to a range between 20 and 24 °C until it arrives at the processing laboratory where it shall be held between 20 and 24 °C until the platelets are separated. The platelet concentrate shall be separated within 4 hours after the collection of the unit of whole blood or plasma.

(c) The time and speed of centrifugation must have been demonstrated to produce an unclumped product, without visible hemolysis, that yields a count of not less than 5.5×10^{10} platelets per unit in at least 75 percent of the units tested.

(d) The volume of original plasma used for resuspension of the platelets shall be determined by the maintenance of a pH of not less than 6.0 during the storage period. The pH shall be measured on a sample of platelets which has been stored for the maximum dating period at the selected storage temperature. One of the following storage temperatures shall be used continuously:

(1) 20 to 24 °C.

(2) 1 to 6 °C.

(e) Final containers used for Platelets shall be colorless and transparent to permit visual inspection of the contents; any closure shall maintain a hermetic seal and prevent contamination of the contents. The container material shall not interact with the contents, under the customary conditions of storage and use, in such a manner as to have an adverse effect upon the safety, purity, potency, or efficacy of the product. At the time of filling, the final container shall be marked or identified